



PATENT ABSTRACTS OF JAPAN

(11) Publication number: **05310625 A**(43) Date of publication of application: **22.11.1993**(51) Int. Cl **C07C 43/13****B01D 15/04, B01D 15/08, B01J 39/06, C07C 41/36**(21) Application number: **04110246**(22) Date of filing: **28.04.1992**(71) Applicant: **MITSUBISHI KASEI CORP**(72) Inventor: **KATSURAGI TOSHIYA
KUSAKABE MASAHIRO****(54) SEPARATION METHOD FOR POLYGLYCERIN
COMPOSITION****(57) Abstract:**

PURPOSE: To easily obtain a polyglycerin composition with desired molecular weight by chromatographic separation of a polyglycerin composition with broad molecular weight distribution using a specific ion exchange resin.

CONSTITUTION: A polyglycerin composition with desired molecular weight can be obtained by simply making a chromatographic separation of (A) a polyglycerin composition with broad molecular weight distribution produced by polymerizing glycerin using (B) a

gel-type strongly acidic cation exchange resin having polystyrene skeleton with a degree of cross-linking of 5-8%. The parent resin for the ion exchange resin is a granular resin with three-dimensional network structure (gel-type) produced by copolymerization using styrene as the chief monomer component and e.g. divinylbenzene as cross-linking monomer. The ion exchange resin to be used has sulfonate group as the strongly acidic cation exchange group at 1.0-2.0meq/l in terms of exchange capacity. The counterion is e.g. H^+ , Na^+ . It is preferable that the original polyglycerin composition contains ≥ 40 wt.% of components 2-8 in degree of polymerization.

COPYRIGHT: (C)1993,JPO&Japio